

Article ID:1005-3085(2011)03-0406-05

Shrinking Projection Methods for Variational Inequalities in Banach Spaces*

GAO Xing-hui¹, ZHOU Hai-yun²

(1- College of Mathematics and Computer Science, Yanan University, Yanan 716000;

2- Department of Mathematics, Shijiazhuang Mechanical Engineering College, Shijiazhuang 050003)

Abstract: In uniformly smooth and uniformly convex Banach spaces, a shrinking projection algorithm is proposed for finding an element of the solution set of variational inequalities, and a strong convergence theorem is proved by using the generalized projection operator, K-K property and other analysis techniques under the conditions of compact mappings weakening continuous mappings. The results of this paper improve and extend recent some relevant results. The proposed algorithm has important applications.

Keywords: shrinking projection method; variational inequalities; K-K property

Classification: AMS(2000) 47H05; 47H09 **CLC number:** O177.91 **Document code:** A

1 Introduction and preliminaries

Throughout this paper, we assume that X is a Banach space and X^* its dual space. We use $\langle \cdot, \cdot \rangle$ to denote the duality pairing between X^* and X . Let C be a subset of X and $T : C \rightarrow X^*$ be a mapping. We consider the following variational inequality problem: find $x \in C$ such that

$$\langle Tx, y - x \rangle \geq 0 \quad (1)$$

for all $y \in C$. A point $x_0 \in C$ is called a solution of the variational inequality (1) if $\langle Tx_0, y - x_0 \rangle \geq 0$ for all $y \in C$. The solution set of the variational inequality (1) is denoted by $VI(C, T)$.

The variational inequality (1) has been extensively investigated^[1-6]. Let X, Y be Banach spaces. A mapping $T : D(T) \subset X \rightarrow Y$ is said to be compact if it is continuous and maps the bounded subsets of $D(T)$ onto the relatively compact subsets of Y , where $D(T)$ is the domain of T . Recently, Fan^[7] proved the following theorem.

Theorem 1 Let X be a uniformly convex and uniformly smooth Banach space and let C be a closed convex subset of X . Suppose that there exists a positive number β such that $\langle Tx, J^{-1}(Jx - \beta Tx) \rangle \geq 0$ for all $x \in C$ and $J - \beta T : C \rightarrow X^*$ is compact. If $\langle Tx, y \rangle \leq 0$ for all $x \in C$ and $y \in VI(C, T)$, then the variational inequality (1) has a solution $x^* \in C$. The sequence $\{x_n\}$ defined by the following iterative scheme

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n \pi_C(Jx_n - \beta Tx_n), \quad n = 1, 2, \dots,$$

Received: 10 July 2009.

Biography: Gao Xinghui (Born in 1975), Female, Master, Associate Professor. Research field: nonlinear functional analysis.

Accepted: 31 Dec 2009.

***Foundation item:** The National Natural Science Foundation of China (10771050); the Scientific Research Program of Shaanxi Provincial Education Department (11JK0486).

where $\{\alpha_n\}$ satisfies $0 < a \leq \alpha_n \leq b < 1$ for all $n \in \mathbb{N}$ and for some positive numbers $a, b \in (0, 1)$ such that $a < b$, \mathbb{N} is the set of positive integers. Then $\{x_n\}$ converges strongly $x^* \in C$.

Question Can the compact mapping $J - \beta T$ in Theorem 1 be weakened to the continuous mapping $J - \beta T$?

The purpose of this paper is to solve the above Question by introducing a new and simple shrinking projection method. The results of this paper improve and extend the corresponding results of Li^[6], Fan^[7] and others.

We denote by J the normalized duality mapping from X to 2^{X^*} defined by

$$Jx = \{f \in X^* : \langle f, x \rangle = \|x\|^2 = \|f\|^2\}, \quad \forall x \in X.$$

Recall that a Banach space X has the K-K property if for any sequence $\{x_n\} \subset X$ that converges weakly to x where also $\|x_n\| \rightarrow \|x\|$, then $\|x_n - x\| \rightarrow 0$ ([8]). It is known that every uniformly convex Banach space has the K-K property.

Let R be the set of real numbers. The functional $V : X^* \times X \rightarrow R$ is defined by

$$V(\phi, x) = \|\phi\|^2 - 2\langle \phi, x \rangle + \|x\|^2,$$

where $\phi \in X^*$ and $x \in X$ ([2]). The functional $V_2 : X \times X \rightarrow R$ is defined by $V_2(x, y) = V(Jx, y)$, for all $x, y \in X$. Let X be a reflexive, strictly convex and smooth Banach space. Then the generalized projection operator $\pi_C : X^* \rightarrow C$ is continuous ([9]). The generalized projection operator π_C and the functional V have the following properties,

- (i) $V : X^* \times X \rightarrow R$ is continuous;
- (ii) $V(\phi, x) = 0$ if and only if $\phi = Jx$;
- (iii) $V(J\pi_C\phi, x) \leq V(\phi, x)$ for all $\phi \in X^*$ and $x \in X$;
- (iv) $\pi_C(Jx) = x$ for all $x \in C$;
- (v) Let X be smooth. For any given $\phi \in X^*$ and $x \in C$, $x \in \pi_C\phi$ if and only if $\langle \phi - Jx, x - y \rangle \geq 0$ for all $y \in C$;
- (vi) The operator $\pi_C : X^* \rightarrow C$ is single-valued if and only if X is strictly convex;
- (vii) Let X be smooth. If $x \in \pi_C\phi$, then $V(Jx, y) \leq V(\phi, y) - V(\phi, x)$, for all $\phi \in X^*$, $y \in C$ ([1, 9]).

Remark 1 If X is a reflexive, strictly convex and smooth Banach space, then for $x, y \in X$, $V_2(x, y) = 0$, i.e., $V(Jx, y) = 0$ if and only if $x = y$.

Lemma 1^[7] Let C be a nonempty closed and convex subset of a reflexive, strictly convex and smooth Banach space X and let $\phi \in X^*$. Then there exists a unique element in C , denoted by $\pi_C\phi$, such that $V(\phi, \pi_C\phi) = \inf_{y \in C} V(\phi, y)$.

Lemma 2^[1] Let X be a reflexive, strictly convex and smooth Banach space with dual space X^* . Let T be an arbitrary operator from X to X^* and let α be an arbitrary fixed positive number. Then the point $x \in C \subset X$ is a solution to the variational inequality (1) if and only if x is a solution of the operator equation in X , $x = \pi_C(Jx - \alpha Tx)$.

Lemma 3^[5] Let X be a uniformly convex Banach space and $B_r(0)$ be a closed ball of X . Then there exists a continuous strictly increasing convex function $g : [0, \infty) \rightarrow [0, \infty)$ with $g(0) = 0$ such that

$$V_2(\alpha x_1 + (1 - \alpha)x_2, y) \leq \alpha V_2(x_1, y) + (1 - \alpha)V_2(x_2, y) - \alpha(1 - \alpha)g(\|x_1 - x_2\|),$$

for all $x_1, x_2, y \in B_r(0)$ and $\alpha \in [0, 1]$.

Lemma 4^[10] Let X be a uniformly convex and smooth Banach space and $\{y_n\}$ and $\{z_n\}$ be two sequences of X . If $V_2(z_n, y_n) \rightarrow 0$ and either $\{y_n\}$ or $\{z_n\}$ is bounded, then $z_n - y_n \rightarrow 0$.

Lemma 5^[4] Let X be a uniformly convex and uniformly smooth Banach space. Then the following inequality holds

$$\|\phi + f\|^2 \leq \|\phi\|^2 + 2\langle \phi, J^{-1}(\phi + f) \rangle, \quad \forall \phi, f \in X^*.$$

Lemma 6 Let X be a uniformly convex and uniformly smooth Banach space and let C be a nonempty closed and convex subset of X . If there exists a positive number β such that

$$\langle Tx, J^{-1}(Jx - \beta Tx) \rangle \geq 0, \quad \forall x \in C, \quad (2)$$

$$\langle Tx, y \rangle \leq 0, \quad \forall x \in C, \quad y \in VI(C, T). \quad (3)$$

Then $VI(C, T)$ is closed and convex.

Proof From the definition of V_2 , the property (iii) of V , (2), (3) and Lemma 4, we can obtain that $VI(C, T)$ is closed and convex.

2 Main results

Theorem 2 Let X be a uniformly convex and uniformly smooth Banach space. Let C be a nonempty closed convex subset of X . Assume that T is an operator of C into X^* which satisfies conditions (2) and (3). Define a sequence $\{x_n\}$ by the following algorithm

$$\begin{cases} x_0 \in X, & C_1 = C, \\ y_n = (1 - \alpha_n)x_n + \alpha_n\pi_C(Jx_n - \beta Tx_n), \\ C_{n+1} = \{z \in C_n : V_2(y_n, z) \leq V_2(x_n, z)\}, \\ x_{n+1} = \pi_{C_{n+1}}Jx_0, \end{cases} \quad (4)$$

where $\{\alpha_n\} \subset (0, 1]$ satisfies $\liminf_{n \rightarrow \infty} \alpha_n > 0$. If $J - \beta T : C \rightarrow X^*$ is continuous, then $\{x_n\}$ converges strongly to $\pi_{VI(C, T)}Jx_0$.

Proof 1) Show that $\pi_{VI(C, T)}Jx_0$ is well defined for every $x_0 \in X$.

From [7], we know that $VI(C, T) \neq \emptyset$. By Lemma 6 and Lemma 1, we have that $\pi_{VI(C, T)}Jx_0$ is well defined.

2) Show that C_n is closed and convex for all $n \in \mathbb{N}$.

This follows from the construction of C_n . We omit the details.

3) Show that $VI(C, T) \subset C_n$ for all $n \in \mathbb{N}$.

It is obvious that $VI(C, T) \subset C = C_1$. Suppose that $VI(C, T) \subset C_n$ for some $n \in \mathbb{N}$. For any $p \in VI(C, T) \subset C_n$, from Lemma 3, we have

$$\begin{aligned} V_2(y_n, p) &= V_2((1 - \alpha_n)x_n + \alpha_n\pi_C(Jx_n - \beta Tx_n), p) \\ &\leq (1 - \alpha_n)V_2(x_n, p) + \alpha_n V_2(\pi_C(Jx_n - \beta Tx_n), p) \\ &\quad - \alpha_n(1 - \alpha_n)g(\|\pi_C(Jx_n - \beta Tx_n) - x_n\|). \end{aligned} \quad (5)$$

From the definition of V_2 , the property (iii) of V , (2), (3) and Lemma 5, we have

$$\begin{aligned} V_2(\pi_C(Jx_n - \beta Tx_n), p) &= V(J\pi_C(Jx_n - \beta Tx_n), p) \\ &\leq \|Jx_n - \beta Tx_n\|^2 - 2\langle Jx_n - \beta Tx_n, p \rangle + \|p\|^2 \\ &\leq \|Jx_n\|^2 - 2\langle Jx_n, p \rangle + \|p\|^2 = V_2(x_n, p). \end{aligned} \quad (6)$$

From (5) and (6), we obtain that $V_2(y_n, p) \leq V_2(x_n, p)$, which implies that $p \in C_{n+1}$ and hence $VI(C, T) \subset C_{n+1}$. Therefore, $VI(C, T) \subset C_n$ for all $n \in \mathbb{N}$.

4) Show that $\lim_{n \rightarrow \infty} V(Jx_0, x_n)$ exists.

In view of (4), we have $x_n = \pi_{C_n} Jx_0$. Since $C_{n+1} \subset C_n$ and $x_{n+1} \in C_{n+1}$ for all $n \in \mathbb{N}$, we have $V(Jx_0, x_n) \leq V(Jx_0, x_{n+1})$. On the other hand, we have from 3) that $V(Jx_0, x_n) \leq V(Jx_0, p)$ for all $p \in VI(C, T)$. It follows that $\lim_{n \rightarrow \infty} V(Jx_0, x_n)$ exists.

5) Show that $x_n \rightarrow p_0 \in C$ as $n \rightarrow \infty$.

From 4), we have $\{x_n\}$ is bounded. Note that X is reflexive, without loss of generality, we can assume that $x_n \rightarrow p_0$ weakly as $n \rightarrow \infty$ (passing to a subsequence if necessary). It is easy to see that $p_0 \in C_n$ for all $n \in \mathbb{N}$. Noticing that $V(Jx_0, x_n) \leq V(Jx_0, x_{n+1}) \leq V(Jx_0, p_0)$, by using the definition of V and weakly lower semi-continuity of $\|\cdot\|^2$, we obtain that

$$V(Jx_0, p_0) \leq \liminf_{n \rightarrow \infty} V(Jx_0, x_n) \leq \limsup_{n \rightarrow \infty} V(Jx_0, x_n) \leq V(Jx_0, p_0).$$

It follows that $V(Jx_0, x_n) \rightarrow V(Jx_0, p_0)$ as $n \rightarrow \infty$. Hence $\|x_n\| \rightarrow \|p_0\|$ as $n \rightarrow \infty$. Since X has the K-K property, we have $x_n \rightarrow p_0$ as $n \rightarrow \infty$.

6) Show that $p_0 \in VI(C, T)$.

Noticing that $x_n = \pi_{C_n} Jx_0$ and $x_{n+1} \in C_{n+1} \subset C_n$, in view of the property (vii), we have

$$V(Jx_n, x_{n+1}) \leq V(Jx_0, x_{n+1}) - V(Jx_0, x_n), \quad \forall n \in \mathbb{N}.$$

From 4), we have

$$\lim_{n \rightarrow \infty} V(Jx_n, x_{n+1}) = \lim_{n \rightarrow \infty} V_2(x_n, x_{n+1}) = 0.$$

Since $V_2(y_n, x_{n+1}) \leq V_2(x_n, x_{n+1})$, we have $\lim_{n \rightarrow \infty} V_2(y_n, x_{n+1}) = 0$. By using Lemma 4, we obtain that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - y_n\| = \lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (7)$$

On the other hand, from (4), we have

$$\begin{aligned} \|x_{n+1} - y_n\| &= \|(1 - \alpha_n)(x_{n+1} - x_n) + \alpha_n(x_{n+1} - \pi_C(Jx_n - \beta Tx_n))\| \\ &\geq \alpha_n \|x_{n+1} - \pi_C(Jx_n - \beta Tx_n)\| - (1 - \alpha_n) \|x_{n+1} - x_n\|. \end{aligned}$$

From the condition $\liminf_{n \rightarrow \infty} \alpha_n > 0$ and (7), we obtain that $\|x_{n+1} - \pi_C(Jx_n - \beta Tx_n)\| \rightarrow 0$ as $n \rightarrow \infty$. Since $x_n \rightarrow p_0$ from 5), we have $\lim_{n \rightarrow \infty} \pi_C(Jx_n - \beta Tx_n) = p_0$. Noticing that π_C and $J - \beta T$ are continuous, we have

$$\lim_{n \rightarrow \infty} \pi_C(Jx_n - \beta Tx_n) = \pi_C(Jp_0 - \beta Tp_0).$$

It follows that $\pi_C(Jp_0 - \beta Tp_0) = p_0$. By Lemma 2, we have $p_0 \in VI(C, T)$.

7) Show that $p_0 = \pi_{VI(C, T)} Jx_0$.

From $x_n = \pi_{C_n} Jx_0$, one sees $\langle Jx_0 - Jx_n, x_n - y \rangle \geq 0$, for all $y \in C_n$. From 3), we have

$$\langle Jx_0 - Jx_n, x_n - w \rangle \geq 0, \quad \forall w \in VI(C, T).$$

Since $J : X \rightarrow X^*$ is demi-continuous, we have $\langle Jx_0 - Jp_0, p_0 - w \rangle \geq 0$, for all $w \in VI(C, T)$.

It follows from the property (v) of π_C that $p_0 = \pi_{VI(C, T)} Jx_0$.

References:

- [1] Alber Y I. Metric and generalized projection operators in Banach spaces: properties and applications[C]// Theory and Applications of Nonlinear Operators of Monotonic and Accretive Type. New York: Marcel Dekker, 1996
- [2] Alber Y I, et al. On the projection methods for fixed point problems[J]. Analysis, 2001, 21: 17-39
- [3] Xu B L. Stable iterative algorithms for general mixed variational inequalities[J]. Chinese Journal of Engineering Mathematics, 2009, 26(1): 151-154
- [4] Chang S S. On Chidume's open questions and approximate solutions of multivalued strongly accretive mapping in Banach spaces[J]. Journal of Mathematical Analysis and Applications, 1997, 216: 94-111
- [5] Chidume C E, Li J L. Projection methods for approximating fixed points of Lipschitz suppressive operators[J]. Panamerican Mathematical Journal, 2005, 15: 29-40
- [6] Li J. On the existence of solutions of variational inequalities in Banach spaces[J]. Journal of Mathematical Analysis and Applications, 2004, 295: 115-126
- [7] Fan J H. A mann type iterative scheme for variational inequalities in noncompact subsets of Banach spaces[J]. Journal of Mathematical Analysis and Applications, 2008, 337: 1041-1047
- [8] Hudzik H, et al. Approximative compactness and full rotundity in Musielak-Orlicz spaces and Lorentz-Orlicz spaces[J]. Zeitschrift für Analysis und ihre Anwendungen, 2006, 25(2): 163-192
- [9] LI J. The generalized projection operator on reflexive Banach spaces and its applications[J]. Journal of Mathematical Analysis and Applications, 2005, 306: 55-71
- [10] Kamimura S, Takahashi W. Strong convergence of a proximal-type algorithm in a Banach space[J]. SIAM Journal on Optimization, 2002, 13: 938-945

Banach 空间中关于变分不等式的收缩投影方法

高兴慧¹, 周海云²

(1- 延安大学数学与计算机科学学院, 延安 716000; 2- 石家庄军械工程学院数学系, 石家庄 050003)

摘 要: 在一致光滑的一致凸的 Banach 空间中, 设计了一种收缩投影算法用以逼近变分不等式的解, 并在紧算子减弱为连续算子的条件下, 利用广义投影算子和 K-K 性质等技巧证明了该算法的强收敛性. 所得结果是近期相关结果的改进与推广, 其算法有重要应用.

关键词: 收缩投影算法; 变分不等式; K-K 性质